

In a nutshell: The Gauss-Seidel method

Given a system of n linear equations in n unknowns $\mathbf{A}\mathbf{u} = \mathbf{v}$, we will use iteration to approximate a solution to this system of linear equations. We will assume that A is either strictly diagonally dominant or symmetric and positive definite, in which case, we are assured that all the diagonal entries are non-zero.

Parameters:

- $\mathcal{E}_{\text{step}}$ The maximum step size allowed before we consider the method to have converged.
- N The maximum number of iterations.

1. Define A_{diag} to be the $n \times n$ matrix of the diagonal entries of A and calculate the inverse A_{diag}^{-1} of this matrix, which is that matrix with the reciprocals of each of the diagonal entries of A_{diag} .
2. Define A_{off} to be the $n \times n$ matrix of the off-diagonal entries of A .
3. Let $\mathbf{u}_0 \leftarrow A_{\text{diag}}^{-1} \mathbf{v}$ and $k \leftarrow 0$.
4. If $k > N$, we have iterated N times, so stop and return signalling a failure to converge.
5. Set $\mathbf{u}_{k+1} \leftarrow \mathbf{u}_k$.
6. For i going from 1 to n ,
 - a. Update the i^{th} entry of $(\mathbf{u}_{k+1})_i \leftarrow (A_{\text{diag}}^{-1})_{i,i} (\mathbf{v} - (A_{\text{off}})_{i,\dots} \mathbf{u}_{k+1})$ where $(A_{\text{off}})_{i,\dots}$ is the i^{th} row of A_{off} .
7. If $\|\mathbf{u}_{k+1} - \mathbf{u}_k\|_2 < \mathcal{E}_{\text{step}}$, return \mathbf{u}_{k+1} .
8. Increment k and return to Step 4.

Note that if A is a sparse matrix (most entries are zero and stored using a sparse-matrix representation), then it is reasonable to calculate $A_{\text{diag}}^{-1} A_{\text{off}}$ first and then replace Step 6a by:

- 6a'. Update the i^{th} entry of $(\mathbf{u}_{k+1})_i \leftarrow \mathbf{u}_{0,i} - (A_{\text{diag}}^{-1} A_{\text{off}})_{i,\dots} \mathbf{u}_{k+1}$ where $(A_{\text{diag}}^{-1} A_{\text{off}})_{i,\dots}$ is the i^{th} row of $A_{\text{diag}}^{-1} A_{\text{off}}$.